



DOI: <https://doi.org/10.29298/rmcf.v11i57.552>

Article

## CO<sub>2</sub> microbiano edáfico en un bosque de *Abies religiosa* (Kunth) Schlttdl. & Cham. en la Ciudad de México

### Edaphic microbial CO<sub>2</sub> in a forest of *Abies religiosa* (Kunth) Schlttdl. & Cham. in Mexico City

Guadalupe Barajas Guzmán<sup>1\*</sup>, Dulce Hernández Rosales<sup>1</sup>, Sally Paredes García<sup>1</sup>, Juan Carlos Peña Becerril<sup>1</sup> y Javier Álvarez Sánchez<sup>1</sup>

#### Resumen

La respiración microbiana tiene un papel primordial en el ciclo del C y contribuye a la producción de CO<sub>2</sub>, por lo que es importante identificar los cambios y agentes que alteran a las comunidades microbianas y, por ende, las tasas de respiración. Pocos estudios se han enfocado en caracterizar la respiración microbiana en los bosques urbanos sujetos a cambios en el uso del suelo. En este estudio se examinó la liberación de CO<sub>2</sub> microbiano del suelo de seis sitios con diferente cobertura arbórea (0, 100, 170, 180, 240 y 310 cm<sup>2</sup>) en un bosque de *Abies religiosa*, y su relación con variables edáficas en la cuenca del río Magdalena en la Ciudad de México. En cada sitio se estableció una parcela de 25 × 25 m, se registró el área basal e identificaron las especies arbustivas y herbáceas; se recolectó suelo para medir el CO<sub>2</sub> potencial liberado por la biomasa bacteriana y fúngica, además se estimó el contenido de humedad, densidad aparente, pH, el Carbono y Nitrógeno totales. La liberación de CO<sub>2</sub> por parte de la biomasa fúngica fue mayor que la bacteriana; en la temporada de lluvias se registraron los valores más altos. Se verificó una relación lineal positiva entre la liberación de CO<sub>2</sub> y el contenido de humedad, la densidad aparente, el pH y la relación C/N. Los suelos en los que hubo menos área basal, liberaron mayor CO<sub>2</sub>. La vegetación y las propiedades del suelo influyeron en la actividad microbiana.

**Palabras clave:** Área basal, biomasa fúngica, bosque urbano, CO<sub>2</sub> microbiano, regeneración natural, relación C/N.

#### Abstract

Microbial respiration plays a key role in the C cycle and contributes to the production of CO<sub>2</sub>, so it is important to identify the changes and agents that alter microbial communities and, therefore, soil respiration rates. Few studies have focused on characterizing microbial respiration in urban forests subject to changes in land use. The production of microbial CO<sub>2</sub> from the soil of six sites with different tree cover in an *Abies religiosa* forest (0, 100, 170, 180, 240 and 310 cm<sup>2</sup>) was examined, and its relation with soil properties in the *Magdalena* river basin in Mexico City. A plot of 25 × 25 m was established in each site, the basimetric area was obtained and the shrub and herbaceous species were identified, soil were collected to measure the CO<sub>2</sub> released by the microbial biomass (bacterial and fungal). The gravimetric moisture content, bulk density, pH, and total C and N were measured too. The CO<sub>2</sub> from the fungal biomass was highest than the bacterial; highest values were observed in the rainy season. A positive linear relationship was found between CO<sub>2</sub> and the moisture content, bulk density, pH and C/N ratio. The soils of sites that are in in early stages of natural regeneration with less basimetric area, produce a high release of CO<sub>2</sub>. Vegetation and soil properties influenced microbial activity.

**Key words:** Basimetric area, fungal biomass, urban forest, microbial CO<sub>2</sub>, natural regeneration, C/N ratio.

Fecha de recepción/Reception date: 4 de junio de 2019

Fecha de aceptación/Acceptance date: 17 de diciembre de 2019

<sup>1</sup>Facultad de Ciencias, Universidad Nacional Autónoma de México. México

\*Autor para correspondencia; correo-e: mgbarajasg@ciencias.unam.mx

## Introduction

The *Magdalena* river basin is part of the conservation soil of Mexico City and is classified as a National Park; despite this, there are changes in land use and areas for grazing and agricultural crops (Jujnovsky *et al.*, 2013), which has negative effects on the physical and chemical properties of the soil, which modify the functioning of the ecosystem (Tischer *et al.*, 2014).

The main plant communities in the basin are the *Pinus hartwegii* Lindl., *Abies religiosa* (Kunth) Schlttdl. & Cham. and *Quercus rugosa* Née forests, with different associations of species within them. Studies in the area have shown that the *Abies religiosa* forest has a high productivity ( $10.78 \pm 1.13 \text{ Mg ha}^{-1} \text{ year}^{-1}$ ) (Delgadillo, 2011); It also records the highest rate of decomposition, since in a previous study, Barajas (2014) concluded that the rate of weight loss of the leaf fraction of *A. religiosa* doubled the value of that of *Q. rugosa* and is higher than that of *P. hartwegii*. In addition, it contains a greater amount of water in the ground and a high quality of organic matter, which favors the activity of the disintegrating organisms (Barajas, 2014). Therefore, it is interesting to evaluate the release of CO<sub>2</sub> in this urban forest in the context of carbon circulation, since there is very little information about it in literature, both for fir and urban forests, as is the case with the *Magdalena* river basin.

On a global scale, soil respiration generates  $68 \text{ to } 77 \times 10^{15} \text{ g C}$  per year (Raich and Schlesinger, 1992); around one third of this C comes from the respiration of the roots and the rest from the activity of microorganisms during the process of decomposition of dead organic matter (Lou and Zhou, 2006; Noh *et al.*, 2016; Galicia *et al.*, 2016), so that this process plays an important role in the global carbon cycle and in the concentration of CO<sub>2</sub> in the atmosphere. Therefore, it is of great importance for its implications on climate change and environmental policies (Latimer and Risk, 2016).

CO<sub>2</sub> production is influenced by several biotic and abiotic factors, as well as by their interaction. Environmental factors include temperature, humidity, O<sub>2</sub> concentration, nutrient availability, soil texture and pH (Wood *et al.*, 2013; Zimmerman *et al.*, 2015). Vegetation affects soil respiration, indirectly, by modifying the microclimate and soil structure, and also by the

quantity and quality of organic materials it provides. Quality is determined by the content of labile and recalcitrant compounds of the litter (C:N ratio); rapid decomposition indicates a higher concentration of labile components in relation to the recalcitrant (C:N low), which promotes the rate of decomposition and, therefore, leads to high respiration rates; and, directly, since the respiration of the roots is one of the main contributors to CO<sub>2</sub> (Hasset and Banwart, 1992; Raich and Tufekcioglu, 2000).


Since all these factors are modified by land use change in, it is interesting to assess their impact on the respiration of the soil biota. Currently, it is estimated that between 30 and 66 % of the national territory are under some agricultural or livestock use (Dupuy *et al.*, 2012) and the loss of forest resources is between 500 000 and 631 000 ha per year (Díaz-Gallegos *et al.*, 2008).

Field studies that estimate respiration in Mexican soils are scarce (García *et al.*, 2001; Campos, 2004; Covalada *et al.*, 2009; Campos, 2014) and laboratory tests (Álvarez and Anzueto, 2004; Ikkonen *et al.*, 2004; Alcántara, 2009; Cruz *et al.*, 2012; Cueva *et al.*, 2016), and none has been carried in the green areas of Mexico City. Therefore, the present work was carried out whose objective was to determine and contrast the microbial activity through breathing, in areas with different basimetric area in the *Abies religiosa* forest of the *Magdalena* river basin.

## **Materials and Methods**

### **Study area**

The study was carried out in the *Abies religiosa* forest belonging to the *Magdalena* River Basin (CRM, for its acronym in Spanish). It is a tall, evergreen forest with dense areas that reaches 100 % of vegetation cover and with the dominance of the tree and shrub strata, although there are also open areas due to disturbances and livestock. It is located on steep slopes, at 2 750-3 500 masl. In Figure 1, the sampling sites along the *Abies religiosa* forest appear in white and the basimetric area corresponding to each site is indicated.




Source: modified from Santibáñez (2009).

*Mapa de altitudes y unidades ambientales de la CRM* = Map of altitudes and environmental units of the CRM; *Zonas de muestreo* = Sampling areas.

**Figure 1.** Magdalena river basin at the southeast of Mexico City.

The minimum rainfall recorded is 1 000 mm in the lower and maximum, and up to 1 500 mm in the highest peaks. There are two types of climates, from 2 400 to 2 800 masl, with a sub-humid climate with an average annual temperature between 12 °C and 18 °C. A semi-cold climate and an average annual temperature between 5 °C and 12 °C predominate from 2 800 to 3 800 masl (García, 1988). Figure 2 shows the temperature and precipitation data corresponding to the study period.





Source: *Desviación Alta al Pedregal* meteorological station.

*Temperatura* = Temperature; *Meses del año* = Months of the year; *Precipitación* = Precipitation.

**Figure 2.** Average temperature and total precipitation per month of the study period in the *Magdalena* river basin of Mexico City.

The soil in the forest of *A. religiosa* in the CRM is a humic Andosol (FAO-UNESCO) with medium silty texture to sandy loam (INEGI, 2006). The canopy reaches 20 to 30 m high; the shrub layer is 2 to 3 m high and a herbaceous layer is present. The following plant associations have been reported: *A. religiosa* - *Roldana angulifolia* (DC.) H. Rob. & Brettell and *A. religiosa* - *Acaena elongata* L. (Almeida *et al.*, 2007).

## Selection of sites

The study was carried out in September 2013 (rainy season) and in May 2014 (dry season). Six sites with different vegetation cover were chosen (Table 1) where a 25 × 25 m square was plotted per site, in which the number of species and the perimeter at breast height (PAP) of *Abies religiosa* trees were recorded to obtain the basimetric area.

There were no trees in the most disturbed sites, the richness of herbs was 18 species and the most abundant was *Bromus carinatus* Hook. & Arn., which is a weed native to Mexico and is used as fodder (Rzedowski and Rzedowski, 2001); at the other end

of the conservation gradient, the conserved site is located with the dominant presence of trees of *A. religiosa*. The site with the greatest richness (26 species) presented a basimetric area of 180 cm<sup>2</sup> and the dominant shrub was *Senecio angulifolius* DC. (Table 1), characteristic species of intermediate successional stages in temperate forests (Rzedowski and Rzedowski, 2001).

**Table 1.** Basimetric area of standing trees, vegetal richness and name of the most abundant species in each one of the sites of the *Abies religiosa* (Kunth) Schltldl.& Cham. forest in the *Magdalena* river basin of Mexico City.

Basimetric area (cm <sup>2</sup> )	0	100	170	180	240	310
Vegetal richness	18	9	24	26	14	2
Dominant species	<i>Bromus carinatus</i> Hook. & Arn.	<i>Acaena elongata</i> L.	<i>Salvia gesneriflora</i> Lindl. & Paxton	<i>Senecio angulifolius</i> DC.	<i>Senecio angulifolius</i> DC.	<i>Abies religiosa</i> (Kunth) Schltldl. & Cham.

In the corner and center of each frame, a soil sample was collected at 10 cm deep with a 200-A Soilmoisture auger (10 cm high × 7 cm in diameter). Five replications per site were obtained in each of the seasons (rainy and dry). The samples were stored in a refrigerator at 4 °C until processing. At the same time, at each point a second unaltered soil sample was collected with a 100 cm<sup>3</sup> auger.

## Preparation and analysis

The first five samples were screened with a 2.38 mm mesh opening to estimate bacterial and fungal respiration, and gravimetric moisture content. In the unaltered sample the apparent density, the concentration of carbon (C) and total nitrogen (N), as well as the pH were measured.

The C and N analyzes were performed in the Soil Fertility Laboratory of the *Colegio de Postgraduados* (Graduate College) (Colpos). Total nitrogen was determined by acid digestion with sulfuric acid and colorimetric analysis (Technicon Industrial Systems, 1977). Total C was determined by the modified Walkley-Black method (Technicon Industrial Systems, 1977).

To determine the apparent density, the unaltered soil sample was dried at 105 °C until it reached a constant weight (Siebe *et al.*, 2006).

The pH was recorded using a PC18 Conductronic potentiometer (Hendershot *et al.*, 2007).

The gravimetric moisture content was determined by weighing 50 g of soil collected in the field and placed in an ED53-UL#02-30777 *Binder* drying oven at 105 °C until reaching a constant weight (Jarrell *et al.*, 1999).

### **Microbial respiration**

The Isermeyer technique (Alef, 1995) was followed to measure the release of CO<sub>2</sub> by the total microbial biomass of the soil, which consists of placing a known weight of soil (not wetted) in a 100 mL beaker inside jars with a tight-fitting lid, in which a 0.05 M NaOH solution was added; these were placed in an incubator at 25 °C for three days. At the end of this period, 5 mL of 0.5 M barium chloride solution (BaCl<sub>2</sub>) and four drops of indicator (0.05 M phenolphthalein solution) were added and the samples turned pink. Finally, the pink solution was titrated with hydrochloric acid (HCl) at a concentration of 0.05 M, until it became colorless. The amount of CO<sub>2</sub> was determined from the volume of HCl used (Alef, 1995).

The same technique was used for fungal biomass, but 85 % lactic acid (C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>) was added to eliminate bacterial populations, and thus, evaluate the release of CO<sub>2</sub> by fungi (Van Netten *et al.*, 1994; Dubal *et al.*, 2004; Wolf *et al.*, 2012). To obtain the release of the bacterial CO<sub>2</sub>, the fungal CO<sub>2</sub> was subtracted from the total microbial CO<sub>2</sub>.

The control bottles had no soil, and with them the same procedure was performed.




## Statistical analysis

For the analysis of the release of CO<sub>2</sub>, an analysis of variance (complying with the assumptions of normality and homocedasticity) was performed in three ways considering as factors the source of CO<sub>2</sub> (bacterial, fungal), the site (basimetric area) and seasonality of rainfall, and when significant differences appeared, the Tukey test was performed. Linear regressions were also made between edaphic variables and CO<sub>2</sub> release. All analyzes were performed with 95 % confidence in the STATISTICA 8.0 statistical package (StatSoft Inc., 2007).

## Results

Significant differences in CO<sub>2</sub> release were recorded between bacterial biomass and fungal biomass ( $F_{(1,96)} = 138.34$ ;  $P < 0.0001$ ). The potential release of CO<sub>2</sub> from the fungal biomass was superior to the bacterial one (Figure 3).




Different letters indicate significant differences ( $P < 0.0001$ ).

*Respiración* = Respiration; *Bacteriana* = Bacterial; *Fúngica* = Fungi.

**Figure 3.** CO<sub>2</sub> potential release of the bacterial and fungi biomass in the *Abies religiosa* (Kunth) Schltdl. & Cham. forest in the *Magdalena* river basin of Mexico City.

The CO<sub>2</sub> release also showed significant differences between seasons ( $F_{(1,96)} = 279.55$ ;  $P < 0.0001$ ). There was 1.6 times greater breathing in the rainy season compared to the dry season (Figure 4).



Different letters indicate significant differences ( $P < 0.0001$ ).

*Respiración* = Respiration; *Lluvias* = Rainy season; *Secas* = Dry season.

**Figure 4.** CO<sub>2</sub> potential release in the rainy season and in the dry season in the *Abies religiosa* (Kunth) Schlttdl. & Cham. forest in the *Magdalena* river basin of Mexico City.


The CO<sub>2</sub> emission at the sites recorded significant differences ( $F_{(5,96)} = 32.4$ ;  $P < 0.0001$ ). The highest value was presented on the site with a basimetric area of 100 cm<sup>2</sup> and the lowest in that with 240 cm<sup>2</sup>; there was a 40 % difference between these two sites (Table 2). At the site with the highest basimetric area (absence of undergrowth and herbaceous stratum), 13 % less CO<sub>2</sub> release occurred compared to the highest value.

**Table 2.** Potential release of microbial CO<sub>2</sub> (mean ± standard deviation) from six sites with different basimetric area in the *Abies religiosa* (Kunth) Schltdl. & Cham. in *Magdalena* river basin of Mexico City.

Basimetric area (cm <sup>2</sup> )	CO <sub>2</sub> (mg CO <sub>2</sub> g <sup>-1</sup> día <sup>-1</sup> ) Average
0	0.106 (±0.055) CD
100	0.155 (±0.072) A
170	0.116 (±0.072) C
180	0.108 (±0.033) CD
240	0.094 (±0.040) E
310	0.135 (±0.042) B

Different letters indicate significant differences ( $P < 0.0001$ ).

The interaction between the source of CO<sub>2</sub> release and the season was significant ( $F_{(1,96)} = 90.42$ ;  $P < 0.0001$ ). The highest value corresponded to the combination of fungal biomass in the rainy season, and the CO<sub>2</sub> release was approximately double than recorded in the dry season in each of the groups of microorganisms (Figure 5).



Different letters indicate significant differences ( $P < 0.0001$ ).

*Respiración* = Respiration; *Lluvias* = Rainy season; *Secas* = Dry season;  
*Bacteriana* = Bacterial; *Fúngica* = Fungi.


**Figure 5.** CO<sub>2</sub> potential release of the bacterial and fungal biomass in the rainy and dry season in the *Magdalena* river basin of Mexico City.

The interaction between the source of CO<sub>2</sub> release and sites with different basimetric area was also significant ( $F_{(5,96)} = 34.94$ ;  $P < 0.0001$ ). The lowest values corresponded to bacterial biomass at sites with 170 and 180 cm<sup>2</sup> of basimetric area, and the highest were at sites of 170 and 310 cm<sup>2</sup> in combination with fungal biomass.

Also, the interaction between season and sites with different basimetric area was significant ( $F_{(5,96)} = 35.18$ ;  $P < 0.0001$ ). In the dry season, the lowest value was found in the site with 240 cm<sup>2</sup> of basimetric area, while the highest corresponded to the site with a basimetric area of 100 cm<sup>2</sup> in the rainy season.

Finally, the interaction between the CO<sub>2</sub> release source, the season and the sites with different basimetric area was significant ( $F_{(5,96)} = 36.65$ ;  $P < 0.0001$ ); the lowest average corresponded to bacterial biomass in the rainy season at the site of 170 cm<sup>2</sup> of basimetric area, and the highest to fungal biomass in rains at the site of 100 cm<sup>2</sup>.


Although differences in CO<sub>2</sub> release between sites were detected, the relationship between the basimetric area and respiration was not significant, but there were linear relationships with some of the soil variables. The potential release of CO<sub>2</sub> from bacterial biomass was linearly and positively related to the gravimetric moisture content in the soil ( $R^2 = 0.36$ ,  $F_{(1,28)} = 15.54$ ;  $P < 0.001$ ) and to the bulk density ( $R^2 = 0.52$ ,  $F_{(1,28)} = 29.89$ ;  $P < 0.0001$ ) in the rainy season (Figure 6).



*Respiración* = Respiration; *Agua en el suelo* = Moisture content; *Densidad aparente* = Apparent density.

**Figure 6.** Relationship between the potential release of CO<sub>2</sub> from bacterial biomass and the gravimetric moisture content (a) and between the apparent density (b) in the rainy season, in the *Magdalena* river basin in Mexico City.

The release of CO<sub>2</sub> by fungal biomass was also linearly and positively related to the gravimetric moisture content in the soil during the dry season ( $R^2 = 0.67$ ,  $F_{(1,28)} = 33$ ;  $P < 0.0001$ ) and with the bulk density of the soil in rains ( $R^2 = 0.55$ ,  $F_{(1,28)} = 33$ ;  $P < 0.0001$ ) (Figure 7).




*Respiración* = Respiration; *Agua en el suelo* = Moisture content; *Densidad aparente* = Apparent density.

**Figure 7.** Relationship between the potential release of CO<sub>2</sub> from fungal biomass and the gravimetric moisture content in the dry season (a) and between the apparent density in the rainy season (b), in the *Magdalena* river basin in Mexico City.

Also in the rainy season, the release of CO<sub>2</sub> by the fungal biomass was linearly and positively related to the pH ( $R^2 = 0.48$ ,  $F_{(1,28)} = 25$ ;  $P < 0.0001$ ) and the quality of the organic matter of the soil (estimated through the C: N ratio) ( $R^2 = 0.44$ ,  $F_{(1,28)} = 22$ ;  $P < 0.0001$ ). The lowest respiration values are located at a pH near 4 and the highest at pH values close to 6; while the activity of fungal biomass increases at high values of the C/N ratio (Figure 8).





*Respiración* = Respiration; *Relación C/N* = C/N ratio.

**Figure 8.** Relationship between the release of CO<sub>2</sub> from fungal biomass and pH (a) and the C / N (b) ratio, in the Magdalena river basin in Mexico City.

## Discussion

The recorded CO<sub>2</sub> values in this work are within the range of values for temperate forests: 0.01 to 0.70 mg CO<sub>2</sub> g<sup>-1</sup> day<sup>-1</sup> (Bailey *et al.*, 2002; Ikkonen *et al.*, 2004; Campos, 2014). Although no relationship was found between the basimetric area and the potential release of CO<sub>2</sub>, the site where a richness of 9 plant species was recorded, with the dominance of *Acaena elongata* and whose basimetric area is 100 cm<sup>2</sup> was the one that recorded the CO<sub>2</sub> highest values, followed by the site where there were only trees (310 cm<sup>2</sup>) of *A. religiosa*, suggesting, as in other studies, that heterotrophic respiration rates are modified according to changes in vegetation structure during succession, which has been observed in both temperate and subtropical forests (Yan *et al.*, 2009; Susyan *et al.*, 2011).

In this study, the release of CO<sub>2</sub> by microbial biomass (bacteria and fungi) was different between sites, but did not follow an ascending pattern depending on the basimetric area or species richness (Table 2). Thus, the highest value of respiration, in addition to registering at a site with few plant species (9), also coincided with the dominance of *Acaena elongata*, a species that indicates disturbance and has a high demand for light, and which tends to grow in spots and dominate the herbaceous stratum (Rzedowski and Rzedowski, 2001), which prevents the establishment of other species.

In two of the sites with significant presence of trees (180 and 240 cm<sup>2</sup> of basimetric area) *Senecio angulifolius*, dominant species that reaches up to 5 m high, favors the entry and establishment of other species in the soil (Rzedowski and Rzedowski, 2001), which means that the sequence is in an intermediate stage; low CO<sub>2</sub> potential release values were recorded here. Instead, Susyan *et al.* (2011) observed that the basal respiration rates of the soil, as well as the microbial biomass increase during the succession process after the abandonment of agricultural land, which is attributed to an increase in the entrances and accumulation of organic C in soil.

In regard to the relationship between CO<sub>2</sub> release and some soil characteristics, a reduction in respiration rates due to an alteration of the soil structure, consistent with human activities within the CRM, was expected; in general, these tend to result in soil compaction, reducing the flow of oxygen and water along the soil profile (Tan, 2000), thus limiting microbial activity, so that the degree of compaction is reflected in high values of apparent density. In this regard, Torbert and Wood (1992) and Pengthamkeerati *et al.* (2005) showed a reduction between 60 and 65 % of soil CO<sub>2</sub> flows with values of apparent density in the range of 1.4 - 1.8 g m<sup>-3</sup>. Additionally, Li *et al.* (2002) observed a decline between 26 and 39 % in the abundance of bacteria, fungi and actinomycetes in the soil as a reaction to an increase in bulk density. In this work the density values were in the range of 0.9 and 1.3 g cm<sup>-3</sup>, which indicates that the soil has no compaction problems and that it has a good structure despite the different land uses that occur in the CRM.

The acid pH and a high content of recalcitrant materials favor the establishment of the fungal communities (Joergensen and Wichern, 2008; Paul, 2015; Kamble *et al.*, 2016), hence a positive and significant linear relationship between the CO<sub>2</sub> release from fungal biomass and these two variables. These results are consistent with those of Lou and Zhou (2006) and Zhang *et al.* (2016), who found that fungi are more resistant to acidic pH than bacteria in forest soils. However, the data in this study show a clear decrease in microbial respiration rates at pH values lower than 5. Sinsabaugh *et al.* (2008) recorded that very low pH values limit the enzymatic extracellular activity of fungi, which would explain what was found in this work.

The C/N ratio of the leaf fraction of the litter of *A. religiosa* is 50 (Barajas, 2014), which indicates that it is a substrate of difficult degradation, while the C/N ratio of soil organic matter in the rainy season goes from 20 to 25 and under dry conditions it reaches only values equal to 15. The highest CO<sub>2</sub> release was recorded at values of 25 in the C/N ratio of the organic matter of the soil, which was observed during the rainy season, so that a good quality, combined with the water present in the soil, favors microbial activity.

Several authors have observed a very marked response of microbial respiration rates in relation to seasonality, presenting the highest values during the rainy season, which is a consequence of a greater availability of nutrients and water in the soil and the consequent activation of the microbial community (Raich and Schlesinger, 1992; Yan *et al.*, 2009; Wood *et al.*, 2013). On the other hand, during periods of drought, access to nutrients becomes limited as the water layer inside the soil is reduced (Lou and Zhou, 2006; Barnard *et al.*, 2013), resulting in lower decomposition rates and therefore breathing. In this work, approximately twice the release of CO<sub>2</sub> was recorded in the rainy season with respect to the dry season, which is consistent with what was reported in studies conducted in temperate ecosystems, in humid tropical and in transformed environments, such as crops and grasslands (Munson *et al.*, 2010; Carbone *et al.*, 2011; Thomey *et al.*, 2011; Galicia *et al.* 2016).

The fact that there has been a positive linear relationship between the release of CO<sub>2</sub> and the moisture content in the soil during the dry season, could be due to the fact that the fungi have greater resistance to desiccation associated with their mycelial growth, which It allows spreading their hyphae along the pores of the soil absorbing nutrients and water (De Vries *et al.*, 2012; Paul, 2015). Likewise, fungi as growth strategies k produce more enzymes to access the use of more organic forms of C accumulated during this period (Waldrop *et al.*, 2006).



## **Conclusions**

The results in this work indicate that the sites in a succession process in their early stages and that have less basimetric area in the CRM, generate a high potential release of CO<sub>2</sub> in the soil.

They also corroborate the relationship between some soil characteristics and microbial activity, the gravimetric moisture content, the pH, the apparent density and the quality of the organic matter modified the potential release of CO<sub>2</sub>. So it is a multifactorial process.

Finally, it is suggested that disturbed areas should be recovered to reduce CO<sub>2</sub> emissions from the soil into the atmosphere of Mexico City.

## **Acknowledgements**

To the Research Program on Climate Change 2012 (PINCC): "The *Magdalena* river basin of Mexico City as a reference site for the monitoring of the effects of climatic change", for the support and sponsoring to accomplish this project.

## **Conflict of interests**

The authors declare no conflict of interests.

## **Contribution by author**

Guadalupe Barajas Guzmán: field work, analysis of results and writing of the article; Dulce Hernández Rosales: field and laboratory work; Sally Paredes García: field and laboratory work; Juan Carlos Peña Becerril: field work and review of the manuscript; Javier Álvarez Sánchez: field work and review of the manuscript.

## References

Alcántara, A. 2009. Producción de CO<sub>2</sub> microbiano en suelos de bosques templados de Oaxaca: una comparación de sitios húmedos y secos. Tesis de Maestría. Instituto de Ecología, Universidad Nacional Autónoma de México. México, D.F., México. 73 p.

Alef, K. 1995. Soil respiration. *In*: Alef, K. and P. Nannipieri (eds.). *Methods in applied soil microbiology and biochemistry*. Academic Press Inc. San Diego, CA, USA. pp. 214-215.

Almeida-Leñero, L., M. Nava, A. Ramos, M. Espinosa, M. J. Ordóñez y J. Jujnovsky. 2007. Servicios ecosistémicos en la cuenca del río Magdalena, Distrito Federal, México. *Gaceta Ecológica* 84-85: 53–64.

Álvarez S., J. and M. Anzueto M. 2004. Soil microbial activity under different corn cropping systems in the highlands of Chiapas, México. *Agrociencia* 38(1): 13-22.

Barajas G., G. 2014. Descomposición, respiración del suelo y macrofauna edáfica en la cuenca del Río Magdalena, México, D. F. *In*: Paz, F. y J. Wong (eds.). *Estado Actual del Conocimiento del Ciclo del Carbono y sus Interacciones en México: Síntesis a 2014*. Serie: Síntesis Nacionales. Programa Mexicano del Carbono. Centro de Investigación y Estudios Avanzados del Instituto Politécnico Nacional, Unidad Mérida. Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco. Texcoco, Edo. de Méx., México. pp. 93-100.

Bailey, V., L. Smith and H. Bolton. 2002. Fungal-to-bacterial ratios in soils investigated for enhanced C sequestration. *Soil Biology & Biochemistry* 34: 997–1007. Doi: 10.1016/S0038-0717(02)00033-0.

Barnard, L., C. Osborne and K. Firestone. 2013. Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. *International Society for microbial Ecology* 7(11): 2229–2241. Doi: 10.1038/ismej.2013.104.

- Campos, C. 2004. Effects of subsistence system on soil surface CO<sub>2</sub>-C flux on Cofre de Perote volcano slopes, Veracruz (Mexico). *Forest Ecology and Management* 199(2-3): 273–282. Doi:10.1016/j.foreco.2004.05.045.
- Campos, C. 2014. Trends in soil respiration on the eastern slope of the Cofre de Perote Volcano (Mexico): Environmental contributions. *Catena* 114: 59-66. Doi:10.1016/j.catena.2013.10.010.
- Carbone, M., C. Still, R. Ambrose, T. Dawson, A. Williams, C. Boot and J. Schimel. 2011. Seasonal and episodic moisture controls on plant and microbial contributions to soil respiration. *Oecologia* 167(1): 265-278. Doi: 10.1007/s00442-011-1975-3.
- Covalada, S., C. Prat, F. García O., J. Etchevers, J. Gallardo. y F. Paz. 2009. Flujos de CO<sub>2</sub> edáfico en un transecto de bosque de pino-encino afectados por actividad antrópica en la microcuenca de Atécuaro (Michoacán, México). *In*: Campo A., J. y M. Conti (eds.). *Emisiones de gases con efecto invernadero en ecosistemas iberoamericanos*. Sociedad Iberoamericana de Física y Química Ambiental. Salamanca, España. pp.123–153.
- Cruz R., E., A. Cruz R., L. Aguilera G., H. Norman M., R. Velázquez, G. Nava B., L. Dendooven y B. Reyes R. 2012. Efecto en las características edáficas de un bosque templado por el cambio de uso de suelo. *Terra Lationamericana* 30(2): 189–197.
- Cueva, A., C. Robles, J. Payan and E. Yépez. 2016. Soil respiration in Mexico: Advances and future directions. *Terra Latinoamericana* 34(3): 253–269. Doi:10.1007/s10021-012-9537-z.
- De Vries, F., M. Liiri, L. Bjornlund, M. Bowker, S. Christensen, H. Setälä and R. Bardgett. 2012. Land use alters the resistance and resilience of soil food webs to drought. *Nature Climate Change* 2: 276–280. Doi: 10.1038/nclimate1368.
- Delgadillo, E. 2011. Productividad primaria neta de los bosques templados de la Cuenca del Río Magdalena. Tesis de Maestría. Facultad de Ciencias, Universidad Nacional Autónoma de México. México, D.F., México. 82 p.

- Díaz-Gallegos, J. R., J. F. Mas y A. Velázquez M. 2008. Monitoreo de los patrones de deforestación en el Corredor Biológico Mesoamericano, México. *Interciencia* 33(12): 882–890.
- Dubal, Z. B., A. M. Paturkar, V. S. Waskar, R. J. Zende, C. Latha, D. B. Rawool and M. M. Kadam. 2004. Effect of food grade organic acids on inoculated *S. aureus*, *L. monocytogenes*, *E. coli* and *S. Typhimurium* in sheep/goat meat stored at refrigeration temperature. *Meat Science* 66 (4): 817-821. Doi: 10.1016/j.meatsci.2003.08.004.
- Dupuy R., J. M., J. L. Hernández S., R. Hernández J., F. Tun D. y F. May P. 2012. Efectos del cambio de uso del suelo en la biomasa y diversidad de plantas leñosas en un paisaje de bosque tropical seco en Yucatán. *Investigación ambiental* 4(1): 130-140.
- Galicia, L., A. Gamboa, S. Cram, B. Chávez, V. Peña V. Saynes y C. Siebe. 2016. Almacén y dinámica del carbono orgánico del suelo en bosques templados de México. *Terra Latinoamericana* 34: 1-29.
- García, E. 1988. Modificaciones al sistema de clasificación climática de Köppen. UNAM. México, D. F., México. 90 p.
- García, N., E. Romero, S. Galicia, A. Ibáñez, B. Platero, R. Ramos and G. Almendros. 2001. Behavior of the soil biogeochemical cycles on control sites of coffee growing agroecosystems from Pluma Hidalgo, Oaxaca (Mexico). *In: Weber, J., E. Jamroz, J. Drozd and A. Karczewska (eds.). Biogeochemical processes and cycling of elements in the environment. Polish Society of Humic Substances. Wroclaw, Poland. pp. 343–344.*
- Hasset, J. and W. Banwart. 1992. *Soils and their environment*. Prentice Hall. Upper Saddle River, NJ, USA. 424 p.
- Hendershot, W., H. Lalonde and M. Duquette. 2007. Soil reaction and exchangeable acidity. *In: Carter, M. R. (eds.). Soil sampling and methods of analysis. Carter CRC Press. Boca Raton, FL, USA. pp. 173–178.*

Ikkonen, E., E. Ángeles-Cervantes y N. E. García-Calderón. 2004. Producción de CO<sub>2</sub> en Andosoles afectados por incendios forestales en el Parque Nacional El Chico, Hidalgo. *Terra Latinoamericana* 22(4): 425-431.

Instituto Nacional de Estadística Geografía e Informática (INEGI). 2006. Cuaderno estadístico delegacional. Magdalena Contreras. INEGI. México, D.F., México. <https://www.inegi.org.mx/app/biblioteca/ficha.html?upc=702825006450> (1 de febrero de 2019).

Jarrell, M., D. Armstrong, F. Grigal, F. Eugene, M. Curtis and H. Wedin. 1999. Soil Water and Temperature Status. *In: Robertson, G. P., D. C. Coleman, C. S. Bledsoe and P. Sollins (eds.). Standard soil methods for long-term ecological research.* Oxford University Press. Oxford, UK. pp. 55-73.

Joergensen, R. and F. Wichern. 2008. Quantitative assessment of the fungal contribution to microbial tissue in soil. *Soil Biology and Biochemistry* 40(12): 2977–2991. Doi: 10.1016/j.soilbio.2008.08.017.

Jujnovsky, J., L. Galván y M. Mazari-Hiriart. 2013. Zonas protectoras forestales: El caso de los bosques de la Cañada de Contreras, Distrito Federal. *Investigación Ambiental* 5(2): 5-15.

Kamble, P. and E. Bååth. 2016. Comparison of fungal and bacterial growth after alleviating induced N-limitation in soil. *Soil Biology and Biochemistry* 103: 97–105. Doi: 10.1016/j.soilbio.2016.08.015.

Latimer, R. and D. Risk. 2016. An inversion approach for determining distribution of production and temperature sensitivity of soil respiration. *Biogeosciences* 13(7): 2111–2122. Doi:10.5194/bg-13-2111-2016.

Li, C. H., B. L. Ma and T. Q. Zhang. 2002. Soil bulk density effects on soil microbial populations and enzyme activities during the growth of maize (*Zea mays* L.) planted in large pots under field exposure. *Canadian Journal of Soil Science* 82(2): 147-154. Doi:10.4141/S01-026.

Luo, Y. and X. Zhou. 2006. Soil respiration and the environment. Academic Press. San Diego, CA, USA. 328 p.

Munson, S., T. Benton, W. Lauenroth and C. Burke. 2010. Soil carbon flux following pulse precipitation events in the shortgrass steppe. *Ecological Research* 25(1): 205-211. Doi:10.1007/s11284-009-0651-0.

Noh, N. J., M. Kuribayashi, T. M. Saitoh, T. Nakaji, M. Nakamura, T. Hiura and H. Muraoka. 2016. Responses of soil, heterotrophic, and autotrophic respiration to experimental open-field soil warming in a cool-temperate deciduous forest. *Ecosystems* 19(3): 504–520. Doi: 10.1007/s10021-015-9948-8.

Paul, E. A. (ed.). 2015. Soil microbiology, ecology and biochemistry. Academic Press. Burlington, MA, USA. 582 p.

Pengthamkeerati, P., P. P. Motavalli, R. J. Kremer and S. H. Anderson. 2005. Soil carbon dioxide efflux from a claypan soil affected by surface compaction and applications of poultry litter. *Agriculture, Ecosystems & Environment* 109 (1-2): 75–86. Doi:10.1016/j.agee.2005.02.013.

Raich, J. and W. Schlesinger. 1992. The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus B: Chemical and Physical Meteorology* 44(2): 81–99. Doi:10.1034/j.1600-0889.1992.t01-1-00001.x.

Raich, J. and A. Tufekcioglu. 2000. Vegetation and soil respiration: Correlations and controls. *Biochemistry* 48(1): 71-90. Doi: 10.1023/A:1006112000616.

Rzedowski, G. y J. Rzedowski. 2001. Flora fanerogámica del Valle de México. Instituto de Ecología, A. C. y Comisión Nacional para el Conocimiento y Uso de la Biodiversidad. México, D. F., México. pp. 33, 38, 235, 936-938.

- Santibáñez A., G. 2009. Composición y estructura del bosque de *Abies religiosa* en función de la heterogeneidad ambiental y determinación de su grado de conservación en la Cuenca del Río Magdalena, México, D.F. Tesis de Maestría. Facultad de Ciencias, Universidad Nacional Autónoma de México. México, D.F., México. 136 p.
- Siebe, C., J. Reinhold y K. Stahr. 2006. Manual para la descripción y evaluación ecológica de suelos en el campo. Sociedad Mexicana de las Ciencias del Suelo. Texcoco, Edo. de Méx., México. 54 p.
- Sinsabaugh, R., C. Lauber, M. Weintraub, B. Ahmed, S. Allison, C. Crenshaw, D. Cusack, S. Frey, M. Gallo, T. Gartner, S. Hobbie, K. Holland, B. Keeler, J. Powers, M. Stursova, C. Takacs-Vesbach, M. Waldrop, M. Wallenstein, D. Zak and H. Zeglin. 2008. Stoichiometry of soil enzyme activity at global scale. *Ecology Letters* 11(11): 1252–1264. Doi:10.1111/j.1461-0248.2008.01245.x.
- StatSoft Inc. 2007. Statistica para windows. Version manual 6.0. StatSoft Inc. Tulsa, OK, USA. n/p.
- Susyan, E., S. Wirth, N. Ananyeva and E. Stolnikova. 2011. Forest succession on abandoned arable soils in European Russia—Impacts on microbial biomass, fungal-bacterial ratio, and basal CO<sub>2</sub> respiration activity. *European Journal of Soil Biology* 47(3): 169–174. Doi: 10.1016/j.ejsobi.2011.04.002.
- Tan, K. 2000. Environmental soil science. Marcel Dekker, Inc. New York, NY, USA. 452 p.
- Technicon Industrial Systems. 1977. Individual/simultaneous determination of nitrogen and/or phosphorus in BD acid digests. Industrial Method No. 329-374. W./B. Technicon Industrial Systems. Tarrytown, NY, USA. pp. 239-374.
- Thomey, M., S. Collins, R. Vargas, J. Johnson, R. Brown, D. Natvig and M. Friggens. 2011. Effect of precipitation variability on net primary production and soil respiration in a Chihuahuan Desert grassland. *Global Change Biology* 17(4): 1505-1515. Doi: 10.1111/j.1365-2486.2010.02363.x.

Tischer, A., K. Potthast and U. Hamer. 2014. Land-use and soil depth affect resource and microbial stoichiometry in a tropical mountain rainforest region of southern Ecuador. *Oecologia* 175(1): 375–393. Doi: 10.1007/s00442-014-2894-x.

Torbert, A. and C. Wood. 1992. Effects of soil compaction and water-filled pore space on soil microbial activity and N losses. *Communications in Soil Science and Plant Analysis* 23(11): 1321–1331. Doi: 10.1080/00103629209368668.

Van Netten, P., J. Huis in't Veld and D. Mossel. 1994. The immediate bactericidal effect of lactic acid on meat-borne pathogens. *Journal of Applied Bacteriology* 77: 490–496. Doi: 10.1111/j.1365-2672.1994.tb04392.x.

Waldrop, M., D. Zak, C. Blackwood, C. Curtis and D. Timan. 2006. Resource availability controls fungal diversity across a plant diversity gradient. *Ecology Letters* 9: 1127–1135. Doi: 10.1111/j.1461-0248.2006.00965.x.

Wolf, M., M. Miller, A. Parks, G. Loneragan, A. Garmyn, L. Thompson, A. Echeverry and M. Brashears. 2012. Validation comparing the effectiveness of a lactic acid dip with a lactic acid spray for reducing *Escherichia coli* O157:H7, *Salmonella*, and Non-O157 Shiga toxigenic *Escherichia coli* on beef trim and ground beef. *Journal of Food Protection* 75(11): 1968 - 1973. Doi: 10.4315/0362-028X.JFP-12-038.

Wood, T., M. Detto and W. Silver. 2013. Sensitivity of soil respiration to variability in soil moisture and temperature in a humid tropical forest. *PLoS ONE* 8(12):1–8. Doi: 10.1371/journal.pone.0080965.

Yan, J., D. Zhang, G. Zhou and J. Liu. 2009. Soil respiration associated with forest succession in subtropical forests in Dinghushan Biosphere Reserve. *Soil Biology y Biochemistry* 41(5): 991–999. Doi: 10.1016/j.soilbio.2008.12.018.

Zhang, Q., J. Wu, F. Yang, Y. Lei, Q. Zhang and X. Cheng. 2016. Alterations in soil microbial community composition and biomass following agricultural land use change. *Scientific Reports* 6: 1-10. Doi: 10.1038/srep36587.

Zimmermann, M., K. Davies, V. Peña and M. Bird. 2015. Impact of temperature and moisture on heterotrophic soil respiration along a moist tropical forest gradient in Australia. *Soil Research* 53(3): 286–297. Doi: 10.1071/SR14217.



All the texts published by **Revista Mexicana de Ciencias Forestales** –with no exception– are distributed under a *Creative Commons* License [Attribution-NonCommercial 4.0 International \(CC BY-NC 4.0\)](https://creativecommons.org/licenses/by-nc/4.0/), which allows third parties to use the publication as long as the work's authorship and its first publication in this journal are mentioned.